

Ana Rita Gomes Teixeira Vaz

Efeito da corticoterapia antenatal na morbilidade e mortalidade de recém-nascidos pré-termo simples e gemelares / Effect of antenatal corticosteroids on morbidity and mortality in preterm singletons and twins

março, 2017

FMUP

Ana Rita Gomes Teixeira Vaz

Efeito da corticoterapia antenatal na morbilidade e mortalidade de recém-nascidos pré-termo simples e gemelares / Effect of antenatal corticosteroids on morbidity and mortality in preterm singletons and twins

Mestrado Integrado em Medicina

Área: Ginecologia e Obstetrícia

Tipologia: Dissertação

Trabalho efetuado sob a Orientação de:

Doutor Nuno Montenegro

E sob a Coorientação de:

Doutora Teresa Rodrigues

Trabalho organizado de acordo com as normas da revista:

Journal of Maternal Fetal and Neonatal Medicine

março, 2017

FMUP

Eu, Ana Rita Gomes Teixeira Vaz, abaixo assinado, nº mecanográfico 201102510, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 1 / 3 / 2017

Assinatura conforme cartão de identificação:

Ana Rita Gomes Teixeira Vaz

NOME

Ana Rita Gomes Teixeira Vaz

NÚMERO DE ESTUDANTE

201102510

DATA DE CONCLUSÃO

01-01-2017

DESIGNAÇÃO DA ÁREA DO PROJECTO

Ginecologia e Obstetrícia

TÍTULO DISSERTAÇÃO/MONOGRÁFIA (riscar o que não interessa)

Effect of antenatal corticosteroids on mortality and morbidity of preterm singletons and twins

ORIENTADOR

Professor Doutor Nuno Montenegro

COORDENADOR (se aplicável)

Professora Doutora Teresa Rodrigues

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA OBRA APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA OBRA (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTA OBRA.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 1 / 3 / 2017

Assinatura conforme cartão de identificação: Ana Rita Gomes Teixeira Vaz

Effect of antenatal corticosteroids on morbidity and mortality in preterm singletons and twins.

Abstract

Purpose: Compare the effect of antenatal corticosteroids (ACS) on neonatal outcomes among singleton and twin pregnancies and the impact of completeness and timing of ministration.

Materials and Methods: Retrospective cohort study involving 951 preterm deliveries (25⁺⁰-34⁺⁶weeks), between 2006 and 2015. Neonatal outcomes were evaluated according to completeness of ACS ("Complete" n=441;"Rescue" n=38;"Incomplete" n=175;"No ACS" n=98) and timing of therapy related to delivery ("Before 7 days" n=260; "After 7 days" n=181).

Results: On Respiratory Distress Syndrome (RDS), odds ratio (OR) for twins was 0.172, 95% confidence interval (CI) 0.047;0.591 and for singletons 0.390 (95%CI 0.214;0.703) for complete or rescue courses, and 0.280 (95%CI 0.069;1.066) for twins and 0.906 (95%CI 0.482;1.698) for singletons for incomplete courses. About the need for mechanical ventilation (MV), twins had OR of 0.189 (95%CI 0.052;0.642) and singletons of 0.404 (95%CI 0.222;0.727) for complete or rescue courses and twins had OR=0.225 (95%CI 0.053;0.874) and singletons of 0.404 (95%CI 0.222;0.727) for incomplete courses. About timing, group "After 7 days" had OR=2.00 for RDS (95%CI 1.21;3.30) and 2.32 (95%CI 1.42;3.78) for MV.

Conclusion: ACS improves neonatal outcomes both in singleton and twins. Delivering seven days after a complete course decreased neonatal morbidity.

Key Words:

Antenatal corticosteroids; twins; preterm; morbidity.

Introduction

Preterm birth is the most costly complication of pregnancy and the leading cause of neonatal morbidity and mortality.^[1,2] There are multiple strategies to minimize the risk and the impact of prematurity, such as ministration of antenatal corticosteroids (ACS), in association with tocolysis, neuroprotection with magnesium sulphate, and neonatal life-saving therapies.^[3] These interventions improve neonatal survival after preterm birth.^[4]

Since Liggins and Howie^[5], numerous investigations have been conducted to ascertain the effect of ACS on prevention of neonatal morbidity and mortality in singleton pregnancies; nowadays, ACS are the cornerstone of prophylactic treatment in preterm birth, between 24⁺⁰ and 34⁺⁶ weeks of gestational age (GA), and its ministration is recommended by the National Institute of Health^[2], the American College of Obstetricians and Gynecologists^[6] and the Royal College of Obstetricians and Gynecologists^[7]. When administered prior to preterm birth, ACS are not only effective in preventing respiratory distress syndrome (RDS) but also in reducing other complications of prematurity, such as intraventricular haemorrhage (IVH), retinopathy of prematurity (ROP), sepsis and necrotizing enterocolitis (NEC), and also neonatal mortality^[3,4]. Authors believe that, in order to achieve maximum effect, the ideal timing of delivery must occur between 24h and seven days after the last dose of therapy.^[1,3,4,6,7]

Despite the significant amount of evidence supporting the impact of ACS in singleton pregnancies between 24 and 34⁺⁶ weeks of GA, there's a substantial lack of information regarding twin pregnancies.^[8] The beneficial effect in singletons has justified ACS use in twin pregnancies, as the mechanism of action is likely to be the same, however the evidence is less robust^[1,4,7,9,10].

We aim to compare the effect of ACS in singleton and twin pregnancies in different neonatal outcomes. We have also tried to investigate the impact of therapeutic completeness (complete, incomplete or rescue) and timing (birth before or after seven days after the last ACS dose).

Materials and Methods

We performed a retrospective cohort study at *Centro Hospitalar de São João*, a tertiary level hospital in Portugal, with average 2500 deliveries per year, preterm (<37 weeks) rate of 10% and very preterm (<32 weeks) of 2%.

Maternal demographic and obstetric characteristics and neonatal outcomes of all singleton and twin pregnant women with preterm deliveries between 25 and 34⁺⁶ weeks from January 2006 to December 2015 ($n=951$) were evaluated. After the approval of the *Ethics Committee* of the hospital, medical records of both mothers and newborns were reviewed separately.

We excluded pregnancies with complications such as twin-to-twin transfusion syndrome ($n=19$), major fetal defects ($n=65$), triplet gestations ($n=5$) and women whose clinical records had important lack of information ($n=25$) or follow up losses ($n=46$). In order to increase internal validity, we eliminated pregnant women with less than 25 weeks of GA on admission ($n=39$) since none was exposed to ACS.

Maternal education, maternal age, body mass index (BMI), smoking habits, alcohol or drug abuse, parity, GA on admission and at the time of delivery, the mode of delivery, chorionicity in twins, *abruptio* placenta, fetal growth restriction (defined by Fenton's growth charts^[11]), fetal Doppler abnormalities, polyhydramnios/oligohydramnios, preterm premature rupture of membranes (PPROM), diabetes, hypertensive, auto immune and thyroid diseases were evaluated.

We ascertained whether the mother received ACS, at which GA, the type of ACS, if the course completeness and the time interval (in days) between the first ministration and delivery, registering the same data for rescue courses. A course of ACS consisted on four 6mg doses of intramuscular dexamethasone at 12h intervals or two 12mg doses of intramuscular betamethasone at 24h interval, as the effect of these two types of ACS is apparently similar^[2]. In our study, the ACS used before February 2014 was betamethasone, and since March 2014, dexamethasone.

For analysis, we divided this population in two major groups: "No ACS" group included pregnant women that were not submitted to antenatal corticosteroids ($n=98$) and the "ACS" group women exposed to this therapy ($n=654$). "ACS" group was divided in three subgroups according to ACS completeness: "Complete" (women that had a complete course of ACS ($n=441$)), "Incomplete" (women submitted to less than the recommended dose ($n=175$)) and "Rescue" (women that had a complete or incomplete course of ACS, followed by another complete course two to three weeks after the first one ($n=38$)).

Our primary outcomes were Respiratory Distress Syndrome (RDS), diagnosed according to the criteria of the Update on the European Consensus Guidelines on the Management of Neonatal Respiratory Distress Syndrome in Preterm Infants (2013) ^[12] and the need for mechanic ventilation (MV). Secondary outcomes were neonatal death (defined as death in the first 28 days of life), admission to the NICU (Neonatal Intensive Care Unit), Apgar Index at the 1st and 5th minute of life, arterial pH value at birth, NEC (defined by the modified Bell staging criteria^[13]), IVH (diagnosed and staged based on Papile classification ^[14]), sepsis, ROP (diagnosed and graded by the International Classification of Retinopathy of Prematurity revisited^[15]) and need for phototherapy. We also performed a composite of neonatal morbidity that included RDS, NEC, ROP, IVH, sepsis and need for admission in the NICU.

The association between ACS and neonatal outcomes was evaluated for singletons and twins and adjusted for GA at delivery, weight of newborn and type of pregnancy (singleton vs. twins). For dichotomous neonatal outcomes, due the smaller sample size, the subgroups "Complete" and "Rescue" were grouped as one.

We also performed a subanalysis with the subgroup "Complete", subsequently divided in class "After 7 days" (if the birth happened more that seven days after the conclusion of the course) and "Before 7 days" (birth in the first seven days after completing the course).As we had only two participants that delivered prior to 24h after the last dose of ACS, we only separated the participants in 2 groups. Results for this analysis are presented in general due to loss of statistical power among twins and are adjusted for GA at delivery and weight of the newborn.

For continuous outcomes we used linear mixed effects analysis with random intercept per birth, when twins were included, and generalized least squares when twins were not included. Linear regression coefficients and the respective 95% confidence interval (95%CI) were used to estimate the association between the exposure and the respective outcome. A final model was estimated that included an interaction term between ACS group and being twin. For categorical outcomes we used unconditional logistic regression to estimate the odds ratios and the respective 95%CI. In twins, the data was aggregated and the outcome had considered when at least one of the twins had the outcome.

Results

In our study, 752 pregnancies were eligible from a total of 951 (79,1%); 199 were excluded by the aforementioned exclusion criteria. As so, 131 twins pregnancies and 621 singleton pregnancies were evaluated: 98 included in the "No ACS" group (76 singleton and 22 twins pregnancies), and 654 on the "ACS" group (545 singleton and 109 twins pregnancies). In this last group, 441 were in the "Complete" subgroup, 175 in the "Incomplete" and 38 in the "Rescue" subgroup. The pregnancies included in the "Complete" subgroup were then categorized according to timing as "After 7 days" (n=181) or "Before 7 days" (n=260).

The groups "No ACS" and "ACS" were similar regarding baseline maternal demographic and obstetric characteristics, except for chorioamnionitis, FGR, fetal Doppler abnormalities, mode of delivery and GA at admission (*table 1*).

Neonatal continuous outcomes according to ACS regimens are shown in *table 2*- in the composite of neonatal morbidity, twins had beta values (β) of -0.226 (95%CI -0.588;0.139) and singletons of -0.312 (95%CI -0.536; -0.088) if exposed to complete courses and β values of 0.044 (95%CI -0.364;0.451) in twins and 0.173 (95%CI 0.072;0.418) in singletons when exposed to incomplete courses.

For dichotomous neonatal outcomes, about need for admission in NICU, in singletons the odds ratio (OR) for complete or rescue courses was 0.887 (95%CI 0.427;1.740) and for incomplete courses 6.217(95%CI 2.212;20.265) and in twins OR for complete or rescue courses was 1.061(95%CI 0.286;3.528) and for incomplete courses 7.381(95%CI 1.046;149.349). Regarding RDS, in singletons OR for complete or rescue courses was 0.390 (95%CI 0.214;0.703) and for incomplete courses 0.906(95%CI 0.482;1.698) and in twins OR for complete or rescue courses was 0.172(95%CI 0.047;0.591) and for incomplete courses 0.280(95%CI 0.069;1.066); about the need for phototherapy, in singletons OR for complete or rescue courses was 1.261 (95%CI 0.865; 1.86) and for incomplete courses 1.515(95%CI 1.010;2.294) and in twins (OR for complete or rescue courses was 0.813(95%CI 0.270;2.287) and for incomplete courses 1.252(95%CI 0.367;4.180). Concerning MV need, in singletons OR for complete or rescue courses was 0.404(95%CI 0.222;0.727) and for incomplete courses 0.819(95%CI 0.435;1.535) and in twins OR for complete or rescue courses was 0.189(95%CI 0.052;0.642) and for incomplete courses 0.225 (95%CI 0.053;0.874). We couldn't find significant differences for the variables neonatal death, NEC, IVH, ROP and sepsis.

Neonatal outcomes according to timing are shown in *table 3*: in RDS, the OR for the group "After 7 days" was 2 (95%CI 1.21;3.30), and in almost all analyzed variables this

group was positively associated with adverse outcomes.

Discussion

Improving pregnancy outcomes is a major goal of healthcare today, and with ACS many premature lives have been saved. The success of this therapy is explained by the fact that synthetic glucocorticoids mimic the developmental maturational changes that normally occur in late gestation in response to rising fetal glucocorticoids.

Despite abundant evidence that use of ACS in singleton pregnant women who are in risk of PTL, between 24 and 34⁺⁶ weeks, has important benefits, data in twin pregnancies the data is limited [1,3,4,7,8-10,16-20].

Our main research question was to determine if clinical neonatal outcomes were equivalent when women received the same therapeutic scheme, independently of plurality.

We started to evaluate maternal characteristics (*table 1*): in 26 parameters, only in five (incidence of chorioamnionitis, fetal growth restriction (FGR), Fetal Doppler abnormalities, mode of delivery and mean GA at admission) there was significant differences between the "No ACS" and the "ACS" group. The higher rate of chorioamnionitis in the "No ACS" group may possibly be explained by less usage of ACS in these patients. FGR and fetal Doppler abnormalities were more common in the "ACS" group possibly because there's a more aggressive treatment in this higher risk groups. Regarding the mode of delivery, the majority of the vaginal preterm births occurred amongst women in the "No ACS" group (in fact, being already in labour was the main reason for not receiving ACS between our participants) with high rates of caesarean delivery in the "ACS" group. Furthermore, we analyzed the frequency of elective and urgent caesareans, and probably because these were the most severe cases of fetal compromise, results showed that the majority of urgent caesareans occurred in the "No ACS" group. Since the association between ACS and neonatal outcomes was not changed after inclusion of mode of delivery in the multivariable model, the final analysis wasn't adjusted for this characteristic.

We analyzed the impact of ACS completeness and timing of delivery. In general, and about the completeness, having a complete or rescue course was better than an incomplete course or no treatment, which may be attributed to inadequate dose or duration of exposure, as it was shown in other studies^[21]. About the impact of different timings between the last ACS dose and delivery, the analysis wasn't able to compare singletons with twins because of our sample size, but the absence of statistical interaction lead us to the presupposition that the effect is similar between this two groups. Delivery in the first seven days after conclusion

of this therapy was significantly associated with lower probability of neonatal death, RDS and need for MV, and despite no statistical significance, other outcomes were compatible, suggesting lower probabilities of developing NEC, need for phototherapy or admission to NICU, and also higher values of umbilical arterial pH, Apgar index at 1st and 5th minute of life and number of days at NICU. Though, the IVH was an exception: our results suggest that newborns delivered more than seven days after ministration had lower rates of this disease, a finding perhaps circumstantial but that can also be explained by the fact the higher GAs at admission are associated with lower prematurity associated complications. Our results in the timing sub analysis are also compatible with literature: Blickstein et al.^[22] found that if the delivery occurred seven days past the ministration, there was no effect in the reduction of RDS, in comparison with delivering in the first seven days, both singleton and twins. Moreover, Gyamfi et al.^[10] agreed by demonstrating that concentrations of betamethasone in cord blood decreased over time, irrespective to plurality.

However, our main research question was to evaluate if results in twins were comparable to singletons. Although in singletons there's a generalized consensus of administration of ACS, twins are treated by extrapolation. In the set of continuous variables, participants were divided in the three subgroups aforesaid ("Complete", "Incomplete" and "Rescue"). In the composite of neonatal morbidity, Apgar indexes and number of days at NICU, there were no differences between singletons and twins, so we believe that the effect of complete courses was equally protective. About the NICU admission, RDS, need for phototherapy and MV, in both singleton and twins doing a rescue or complete course was better than doing an incomplete one, which was also better than no treatment at all. When evaluating the difference of impact of ACS completeness in twins versus singletons ^[23], we found that concerning admission in NICU, the association was 20% lower, so we believe that effect is similar in singletons and twins. In respect of RDS, as the association was 43% higher in singletons, we report that when twin pregnancies are submitted to complete regimens, they have less risk of developing this condition, in comparison with singletons. Also, about the need for MV and phototherapy, the difference was of 46% and 67%, respectively, with higher protection in twin pregnancies. So, our results suggest that despite ACS protection both in singletons and twins, on respect of respiratory outcomes, twins may possibly be more protected.

Authors suggest that ACS ministration doesn't reduce the incidence of RDS in preterm twins as it does in singletons, so that there's an interaction with plurality ^[1, 3,16-18]. Because the recommended dose of ACS is administered irrespective of maternal mass or fetal number, it has been hypothesized that the supposed suboptimal benefits of ACS

treatment in twin pregnancies may be attributable to greater degrees of maternal physiological changes in twin pregnancies, such as greater maternal blood volume expansion and different clearance of betamethasone^[24].

However, this hypothesis was challenged by recent studies that demonstrated no difference between either the pharmacokinetics (clearance and volume of distribution) or the maternal serum or cord blood concentrations between singletons and twins^[10,19]. Other studies report that the effect of ACS is present in twin pregnancies, so that the actual dosage and timing is effective ^[1,8,9,19,20]. Recently, Salim et al. demonstrated that serum betamethasone concentration is measurable with ELISA kit, paving the way for future investigations to determine the optimum concentration that would be clinically^[25].

Our results are in this perspective too: apparently the number of fetus isn't a determinant of the effect of this therapy in the majority of our evaluated outcomes, so that giving this therapy to singleton or twin pregnant women seems to produce the same effects. We also highlight that, in the respiratory outcomes, when exposed to the same therapeutic scheme, twins were even more protected. However, there's still need to be sure which dosage of ACS is effective not only in singletons but mostly in twin pregnancies.

Our study had several strengths: the inclusion criteria were broad, and the analysis did not exclude fetal abnormalities compatible with life. In contrast to other studies examining neonatal outcomes, we included detailed antenatal and pregnancy characteristics. Though, the inherent weaknesses of a cohort study design should be recognized: the retrospective nature invites the possibility of sampling bias leading to differences in major adverse outcome measures. Also, the number of patients was not large enough to allow us to have statistically significant results in all of our outcomes, which is not unique to our study.

Although we think it would be important to differentiate between mono and dichorionic twin pregnancies, we did not have enough sample size to perform this analysis. Further observational studies in large cohorts of twins are warranted to increase the power to find differences in neonatal outcomes, as well as increase our confidence in the safety of using this dose and timing of ACS in twin pregnancies, but we highlight the possible ethical issues that may rise with the omission of therapy to the "control" group. A large, randomized, prospective trial would evaluate the difference of impact of the corticotherapy in singleton and twin pregnancy.-

In conclusion, there were differences in the effect of ACS between singletons and twins in RDS, need for MV and phototherapy.

Acknowledgments

We thank José Bernardo for his contribution on reviewing the writing of this paper.

Disclosure Statement

The authors report no conflicts of interest.

REFERENCES

- [1] Kuk JY, An JJ, Cha HH, Choi SJ, Vargas JE, Oh SY, Roh CR, Kim JH. Optimal time interval between a single course of antenatal corticosteroids and delivery for reduction of respiratory distress syndrome in preterm twins. *Am J Obstet Gynecol*. 2013 Sep;209(3):256.e1-7.
- [2] Antenatal Corticosteroids Revisited: Repeat Courses. NIH Consens Statement Online 2000 August 17-18; [cited year, month, day]; 17(2): 1-10.
- [3] Choi SJ, Song SE, Seo ES, Oh SY, Kim JH, Roh CR. The effect of single or multiple courses of antenatal corticosteroid therapy on neonatal respiratory distress syndrome in singleton versus twin pregnancies. *Aust N Z J Obstet Gynaecol*. 2009 Apr;49(2):173-9.
- [4] Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev*. 2006 Jul 19;(3):CD004454. Review.
- [5] Liggins GC, Howie RN. A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory distress syndrome in premature infants. *Pediatrics*. 1972 Oct;50(4):515-25
- [6] American College of Obstetricians and Gynecologists.; Society for Maternal-Fetal Medicine.. ACOG Practice Bulletin No. 144: Multifetal gestations: twin, triplet, and higher-order multifetal pregnancies. *Obstet Gynecol*. 2014 May;123(5):1118-32
- [7] Royal College of Obstetricians and Gynaecologists. Antenatal corticosteroids to reduce neonatal morbidity. Green-top Guideline, vol 7. London: RCOG; 2010
- [8] Melamed N, Shah J, Yoon EW, Pelausa E, Lee SK, Shah PS, Murphy KE; Canadian Neonatal Network Investigators.. The role of antenatal corticosteroids in twin pregnancies complicated by preterm birth. *Am J Obstet Gynecol*. 2016 Oct;215(4):482.e1-9.
- [9] Battista L, Winovitch KC, Rumney PJ, Davis E, Hagemann C, Wing DA. A case-control comparison of the effectiveness of betamethasone to prevent neonatal morbidity and mortality in preterm twin and singleton pregnancies. *Am J Perinatol*. 2008 Aug;25(7):449-53.
- [10] Gyamfi C, Mele L, Wapner RJ, et al. The effect of plurality and obesity on betamethasone concentrations in women at risk for preterm delivery. *Am J Obstet Gynecol*. 2010 Sep;203(3):219.e1-5.
- [11] Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr*. 2013 Apr 20;13:59. doi: 10.1186/1471-2431-13-59
- [12] Sweet DG, Carnielli V, Greisen G, et al. European consensus guidelines on the management of neonatal respiratory distress syndrome in preterm infants--2013 update. *Neonatology*. 2013;103(4):353-68.
- [13] Walsh MC, Kliegman RM. Necrotizing enterocolitis: treatment based on staging criteria. *Pediatr Clin North Am*. 1986 Feb;33(1):179-201.
- [14] Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal

- and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr*. 1978 Apr;92(4):529-34.
- [15] International Committee for the Classification of Retinopathy of Prematurity.. The International Classification of Retinopathy of Prematurity revisited. *Arch Ophthalmol*. 2005 Jul;123(7):991-9.
- [16] Quist-Therson EC, Myhr TL, Ohlsson A. Antenatal steroids to prevent respiratory distress syndrome: multiple gestation as an effect modifier. *Acta Obstet Gynecol Scand*. 1999 May;78(5):388-92.
- [17] Viteri OA, Blackwell SC, Chauhan SP, et al. Antenatal Corticosteroids for the Prevention of Respiratory Distress Syndrome in Premature Twins. *Obstet Gynecol*. 2016 Sep;128(3):583-91.
- [18] Blickstein I, Shinwell ES, Lusk A, et al. Plurality-dependent risk of respiratory distress syndrome among very-low-birth-weight infants and antepartum corticosteroid treatment. *Am J Obstet Gynecol*. 2005 Feb;192(2):360-4
- [19] Della Torre M, Hibbard JU, Jeong H, Fischer JH. Betamethasone in pregnancy: influence of maternal body weight and multiple gestation on pharmacokinetics. *Am J Obstet Gynecol*. 2010 Sep;203(3):254.e1-12.
- [20] Boghossian NS, McDonald SA, Bell EF, et al. Association of Antenatal Corticosteroids With Mortality, Morbidity, and Neurodevelopmental Outcomes in Extremely Preterm Multiple Gestation Infants. *JAMA Pediatr*. 2016 Jun 1;170(6):593-601.
- [21] Elimian A, Figueroa R, Spitzer AR, Ogburn PL, Wiencek V, Quirk JG. Antenatal corticosteroids: are incomplete courses beneficial? *Obstet Gynecol*. 2003 Aug;102(2):352-5.
- [22] Blickstein I. Antenatal corticosteroids: current controversies. *J Perinat Med*. 2016 Apr 6.
- [23] Maldonado G, Greenland S. Simulation study of confounder-selection strategies. *Am J Epidemiol*. 1993 Dec 1;138(11):923-36
- [24] Ballabh P, Lo ES, Kumari J, Cooper TB, Zervoudakis I, Auld PA, Krauss AN. Pharmacokinetics of betamethasone in twin and singleton pregnancy. *Clin Pharmacol Ther*. 2002 Jan;71(1):39-45.
- [25] Salim R, Suleiman A, Colodner R, Nachum Z, Goldstein LH, Shalev E. Measurement of betamethasone concentration in maternal serum treated for fetal lung maturity; Is it feasible? *Reprod Biol Endocrinol*. 2016 Feb 10;14:7

Table 1: Maternal Demographic and Obstetric Characteristics.

	TOTAL	NO ACS (n (%))	ACS (n (%))	p value
Years of education				0.931
<12 years	276	34 (12.3)	242 (87.7)	
> 12 years	432	51 (11.8)	381 (88.2)	
Tobacco				0.867
Yes	50	7 (14.0)	43 (86.0)	
Alcohol				0.379
Yes	6	2 (33.3)	4 (66.7)	
Drugs				0.270
Yes	1	1 (100)	0 (0)	
Parity				0.927
Nulliparous	361	46 (12.7)	315 (87.3)	
Multiparous	393	52 (13.2)	341 (86.8)	
Conception				0.255
Not Spontaneous	75	6 (8.0)	69 (92.0)	
Spontaneous	665	89 (13.4)	576 (86.6)	
DM				0.788
Yes	87	10 (11.5)	77 (88.5)	
Hypertensive Disease				0.383
Yes	176	19 (10.8)	157 (89.2)	
Mode of Delivery				<0.001
Vaginal	37	11 (29.7)	26 (70.3)	
Urgent Cesarean	323	44 (13.6)	279 (86.4)	
Eletive Cesarean	123	7 (5.70)	116(94.3)	
Polihydramnios/Oligohydramnios				0.102
Yes	115	9(7.8)	106(92.2)	
PPROM				0.346
Yes	253	28 (11.1)	225 (88.9)	
Abruptio Placenta				0.093
Yes	39	9 (23.1)	30 (76.9)	
FGR				0.014
Yes	169	12(7.1)	157(92.9)	
DFA				0.005

Yes	113	5 (4.4)	108(95.6)	
Thyroid pathology				0.879
Yes	37	4 (10.8)	33 (89.2)	
Chorioamnionitis				0.036
Yes	6	3 (50.0)	3 (50.0)	
Auto-Immune Diseases				0.819
Yes	22	2 (9.1)	20 (90.9)	
	Mean (SD)	Mean (SD)	Mean (SD)	
Maternal Age	31.03 (5.91)	31.60 (6.436)	30.94 (5.833)	0.931
Body Mass Index	26.57 (5.299)	27.57 (5.84)	26.438 (5.21)	0.115
GA at admission	32.17 (2.56)	33.10 (2.38)	32.03 (2.56)	<0.001

DFA: Doppler blood flow abnormalities; DM: Diabetes Mellitus; FGR: Fetal Growth Restriction; GA: Gestational Age

Table 2: Impact of different ACS regimens in neonatal outcomes represented by continuous variables

	TOTAL ¹	TWINS	SINGLETON	p interaction
	β (95% CI)	β (95% CI)	β (95% CI)	
Apgar 1				0.760
No ACS (Ref)	Ref	Ref	Ref	
Complete	0.833(0.430;1.23)	1.18 (0.544;1.81)	0.707 (0.210;1.20)	
Incomplete	0.289 (-0.160;0.737)	0.289 (-0.434;1.01)	0.269 (-0.280;0.819)	
Rescue	1.419(0.737;2.10)	1.81(0.739;2.886)	1.30(0.467;2.15)	
Apgar 5				0.428
No ACS(Ref)	Ref	Ref	Ref	
Complete	0.627 (0.292;0.960)	0.759 (0.272;1.246)	0.573 (0.150;0.996)	
Incomplete	0.385 (0.014;0.756)	0.041 (-0.512;0.595)	0.489 (0.022;0.955)	
Rescue	1.214(0.651;1.77)	1.21(0.385;2.03)	1.237(0.522;1.951)	
pH				0.657
No ACS(Ref)	Ref	Ref	Ref	
Complete	0.051(0.014;0.090)	0.0248 (-0.023;0.0734)	0.060(0.013;0.107)	
Incomplete	0.020 (-0.022;0.062)	-0.0245 (-0.077;0.0286)	0.034 (-0.079;0.085)	
Rescue	0.059(0.003;0.115)	0.0367(-0.036;0.101)	0.061(-0.008;0.130)	
Days at NICU				0.667
No ACS(Ref)	Ref	Ref	Ref	
Complete	-3.19 (-7.81;1.44)	-2.44 (-9.94;5.052)	-2.98 (-8.61;2.64)	
Incomplete	0.484 (-4.67;5.63)	-1.62 (-10.22;6.971)	1.47 (-4.74;7.69)	
Rescue	-0.916(-8.74;6.91)	6.99(-5.73;19.73)	-2.54 (-12.07;6.97)	
Composite				0.800
No ACS	Ref	Ref	Ref	
Complete	-0.301 (-0.492; -0.110)	-0.224 (-0.588; 0.139)	-0.312 (-0.536 ; -0.088)	
Incomplete	0.142 (-0.067; 0.352)	0.044 (-0.364 ; 0.451)	0.173 (-0.072 ; 0.418)	
Rescue	-0.297 (-0.596; 0.036)	-0.166 (-0.766;0.434)	-0.298 (-0.670;0.074)	

β : beta-value; 95%CI: 95% Confidence Interval NICU: Neonatal Intensive Care Unit; Apgar1: Apgar Index at 1st minute of life; Apgar5: Apgar index at the 5th minute of life; Composite: Included RDS, NEC, IVH, ROP, sepsis and need for admission in the NICU.

¹ Adjusted for type of pregnancy (singleton or twins), gestational age at delivery and weight of the newborn.

Table 3: Impact of different ACS timings in neonatal outcomes

β (95% CI)	
Apgar 1	
Before 7 days	Ref
After 7 days	-0.057 (-0.410;0.295)
Apgar 5	
Before 7 days	Ref
After 7 days	-0.042 (-0.351;0.266)
pH	
Before 7 days	Ref
After 7 days	-0.025 (-0.055;0.0055)
Days at NICU	
Before 7 days	Ref
After 7 days	0.826 (-3.332 ; 4.98)
OR (95% CI)	
NICU	
Before 7 days	Ref
After 7 days	1.093 (0.631 ; 1.894)
RDS	
Before 7 days	Ref
After 7 days	2.00 (1.21;3.30)
NEC	
Before 7 days	Ref
After 7 days	1.56 (0.56;4.34)
Photo	
Before 7 days	Ref
After 7 days	1.075 (0.68 ; 1.70)
IVH	
Before 7 days	Ref
After 7 days	0.673 (0.233;1.94)
Death	
Before 7 days	Ref
After 7 days	2.20 (1.016;4.77)
MV	
Before 7 days	Ref
After 7 days	2.32 (1.42;3.78)

β: beta-value; **OR:** Odds Ratio; **95%CI:** 95% Confidence Interval; **NICU:** Neonatal Intensive Care Unit; **Apgar1:** Apgar Index at 1st minute of life; **Apgar5:** Apgar index at the 5th minute of life; **RDS:** Respiratory distress syndrome; **NEC:** Necrotizing enterocolitis; **Photo:** phototherapy; **IVH:** Intraventricular hemorrhage; **MV:** Mechanical ventilation. Adjusted for type of pregnancy (singleton or twins) and weight of the newborn.

ANEXO 1 - Regras da Revista Journal of Maternal, Fetal & Neonatal Medicine

The Journal of Maternal-Fetal & Neonatal Medicine publishes the following types of articles: The space available in the printed journal is restricted and authors should follow the below word limits, otherwise this may result in their submission being un-submitted or delayed.

Original Articles: The maximum length is 3000 words (excluding references), including headings and 200-word abstract, maximum of 3 figures and/or tables and up to 30 references.

Review articles: Review articles should examine published research on topics relevant to maternal-fetal medicine. The review article should provide a critical analysis of the available information, should lead to a rational conclusion, and highlight areas of future investigation. The maximum length is 3000 words (excluding references), including headings and 200-word abstract, maximum of 3 figures and/or tables, and up to 30 references.

Short Reports: These should be of original laboratory or clinical contributions. The maximum length is 1500 words (excluding references), including headings and 100-word abstract, maximum of 1 figure and/or table, and up to 10 references.

Letters to the Editor: Letters to the Editor may offer criticism of published material in an objective, constructive and educational manner. Within these limits, Letters to the Editor may be provocative and inductive of further debate. They may also discuss matters of general interest. The material for such can be taken from any source of information so long as it pertains to the general field of Maternal-Fetal Medicine, Newborn Medicine, Perinatal Genetics, and Perinatal Ethics in the broadest sense. They will be reviewed by the appropriate editor and will be subject to editing and possible abridgement. If accepted, a copy will be sent to the author(s) of the original article referred to in the Letter to the Editor, giving the author(s) the opportunity to provide a rebuttal with new material considered for publication with the Letter to the Editor.

Opinions and Hypotheses: These should be 400-600 words in length with one figure or table and a maximum of five references.

Education and Debate Articles: These are usually invited of maximum 2000 words, but reports on all aspects of medicine and health are welcomed. They will be peer-reviewed, and should contain an unstructured abstract of no more than 150 words.

General Guidelines

Please write clearly and concisely, stating your objectives clearly and defining your terms. Your arguments should be substantiated with well reasoned supporting evidence.

In writing your paper, you are encouraged to review articles in the area you are addressing which have been previously published in the Journal, and where you feel appropriate, to reference them. This will enhance context, coherence, and continuity for our readers.

Authors should include telephone and fax numbers as well as e-mail addresses on the cover page of manuscripts.

Articles should be written in English. Authors whose native language is not English are requested to have their manuscript checked for linguistic correctness before submission. For a list of resources we recommend for language editing please click [here](#).

Submissions should include, where appropriate, a formal statement that ethical consent for the work to be carried out has been given.

Submission Guidelines:

All submissions should be made online at The Journal of Maternal-Fetal & Neonatal Medicine's ScholarOne Manuscripts site: <http://mc.manuscriptcentral.com/djmf>

New users should first create an account. Once a user is logged onto the site, submissions should be made via the Author Centre. If you are a new author but have previously reviewed for the Journal through the online system, you should use your existing reviewer username and password to log in as an author.

If you experience any problems with your submission or with the site, please contact ScholarOne support through the 'get help now' link.

Full completion of the "Add a Reviewer" section is required as this may speed up the review process in cases where reviewers are difficult to locate or unavailable. Please add the contact details for 3 preferred reviewers. Please do not suggest anyone from the same department as any of the authors.

Ethical Review

The Journal supports the principles of the Declaration of Helsinki, and expect that the authors of papers submitted to the Journal will have obtained ethical consent and followed those legal and regulatory requirements for human experimentation with drugs, including informed consent, according to procedures which apply in their institution and country. The Code of Ethics of the World Medical Association (Declaration of Helsinki) represents a minimal requirement. Please click on Ethics and Consent for further information.

File preparation and types

Manuscripts are preferred in Microsoft Word format (.doc files). Documents must be double-spaced, with margins of one inch on all sides. Tables and figures should not appear in the main text, but should be uploaded as separate files and designated with the appropriate file type upon submission. References should be given in Council of Science Editors (CSE) Citation & Sequence format (see References section for examples).

Structure of Paper

Manuscripts should be structured into headed sections as follows: Title page, Abstract, Introduction, Methods, Results, Discussion, Acknowledgements, Declaration of Interest statement, References, Tables and Figures. Each section should begin on a new sheet and be identified with the shoulder heading. Other subsection headings within the main headings may be used but should be limited.

Title Page

A title page should be provided comprising the manuscript title plus the full names and affiliations of all authors involved in the preparation of the manuscript. One author should be clearly designated as the corresponding author and full contact information, including phone number and email address, provided for this person. A short title (no more than 20 letters) and Keywords (5-8) that are not in the title should also be included on the title page. The keywords will assist indexers in cross indexing your article.

Abstract

An abstract not exceeding 200 words should state the aim of the study, the main findings, and how the results were interpreted. Abstracts for Short Reports should not exceed 100 words.

Instructions for preparing structured abstracts:

Structured abstracts should be no more than 200 words and consist of four paragraphs under the headings:

Objective A precise statement of the primary objectives of the study, including the primary focus (e.g. diagnosis, prognosis, prevention) and information concerning the specific population, test, or outcome being discussed.

Methods How the study was performed, including details of clinical and/or technical procedures.

Results The salient results of the study.

Conclusions The conclusions and their clinical application; the need for new studies may be suggested. Equal emphasis should be given to positive and negative findings of equal scientific merit.

Style

Define abbreviations when they first occur in the manuscript and from there on use only the abbreviation. When many unusual abbreviations are used, list them alphabetically with their definitions on a separate page. Whenever standardized abbreviations are available, use those; create new abbreviations only if absolutely unavoidable. Use generic names for drugs.

If identification of a brand name is wanted, insert it in parentheses together with the manufacturer's name and address after the first mention of the generic name, noting trademark (TM) or registered (®) if applicable. Scientific measurements should be given in SI units, except for blood pressure, which should be expressed in mmHg.

Clinical Trials

Where applicable, authors reporting phase II and phase III randomized controlled trials should refer to the CONSORT Statement (www.consort-statement.org) for recommendations to facilitate the complete and transparent reporting of trial findings. Reports that do not conform to the CONSORT guidelines may need to be revised before formal review.

For our Clinical Trials Registry policy please see:

http://informahealthcare.com/userimages/ContentEditor/1255620275152/Clinical_Trials_Registry.pdf

Acknowledgments and Declaration of Interests

Acknowledgments and Declaration of interest sections are different, and each has a specific purpose.

The Acknowledgments section details special thanks, personal assistance, and dedications.

Contributions from individuals who do not qualify for authorship should also be acknowledged here.

Declarations of interest, however, refer to statements of financial support and/or statements of potential conflict of interest. Within this section also belongs disclosure of scientific writing assistance (use of an agency or agency/ freelance writer), grant support and numbers, and statements of employment, if applicable.

Acknowledgments section

Any acknowledgments authors wish to make should be included in a separate headed section at the end of the manuscript preceding any appendices, and before the references section. Please do not incorporate acknowledgments into notes or biographical notes.

Declaration of Interest section

All declarations of interest must be outlined under the subheading "Declaration of interest". If authors have no declarations of interest to report, this must be explicitly stated. The suggested, but not mandatory, wording in such an instance is: The authors report no declarations of interest. When submitting a paper via ScholarOne Manuscripts, the "Declaration of interest" field is compulsory (authors must either state the disclosures or report that there are none). If this section is left empty authors will not be able to progress with the submission.

Please note: for NIH/Wellcome-funded papers, the grant number(s) must be included in the Declaration of Interest statement.

[Click here to view our full Declaration of Interest Policy.](#)

References

References should follow the Council of Science Editors (CSE) Citation & Sequence format. Only works actually cited in the text should be included in the references. Indicate in the text with Arabic numbers inside square brackets. Spelling in the reference list should follow the original. References should then be listed in numerical order at the end of the article. Examples are provided as follows:

Journal article: [1] Steiner U, Klein J, Eiser E, Budkowski A, Fetters LJ. Complete wetting from polymer mixtures. *Science* 1992;258:1122-1129.

Book chapter: [2] Kuret JA, Murad F. Adenohypophyseal hormones and related substances. In: Gilman AG, Rall TW, Nies AS, Taylor P, editors. *The pharmacological basis of therapeutics*. 8th ed. New York: Pergamon; 1990. pp 1334-1360.

Conference proceedings: [3] Irvin AD, Cunningham MP, Young As, editors. *Advances in the control of Theileriosis*. International Conference held at the International Laboratory for Research on Animal Diseases; 1981Feb 9-13; Nairobi. Boston: Martinus Nijhoff Publishers; 1981.427p.

Dissertation or Thesis: [4] Mangie ED. A comparative study of the perceptions of illness in New Kingdom Egypt and Mesopotamia of the early first millennium [dissertation]. Akron (OH): University of Akron; 1991. 160 p. Available from: University Microfilms, Ann Arbor MI; AAG9203425.

Journal article on internet: [5] Loker WM. "Campesinos" and the crisis of modernization in Latin America. *Jour of Pol Ecol* [serial online] 1996; 3(1). Available: http://www.library.arizona.edu/ej/jpe/volume_3/ascii-lokeriso.txt via the INTERNET. Accessed 1996 Aug 11.

Webpage: [6] British Medical Journal [Internet]. Stanford, CA: Stanford Univ; 2004 July 10 - [cited 2004 Aug 12]; Available from <http://bmj.bmjournals.com/>

Internet databases: [7] Prevention News Update Database [Internet]. Rockville (MD): Centers for Disease Control and Prevention (US), National Prevention Information Network. 1988 Jun - [cited 2001 Apr 12]. Available from <http://www.cdcnpin.org/>

Further examples and information can be found in *The CSE Manual for Authors, Editors, and Publishers*, Seventh Edition. Periodical abbreviations should follow the style given by Index Medicus.

Tables and Figures

The number of tables and figures should be kept to a minimum. Repetition/overlap of information given in tables/figures/text should be avoided as possible. The place at which a table or figure is to be inserted in the printed text should be indicated clearly on the manuscript.

Each table and/or figure must have a legend that explains its purpose without reference to the text. Each table and/or figure must be uploaded separately from the main document. Charts and tables are considered textual and should also be supplied in a format compatible with MS Word.

Tables: should be used only when they can present information more efficiently than running text.

Care should be taken to avoid any arrangement that unduly increases the depth of a table, and the column heads should be made as brief as possible, using abbreviations liberally. Tables should be submitted on separate sheets, numbered in Roman numerals, and their position indicated in the text (e.g., Table I). Each table should have a short, self-explanatory title. Vertical rules should not be used to separate columns. Units should appear in parentheses in the column heading but not in the body of the table. Any explanatory notes should be given as a footnote at the bottom of the table.

Figures : All illustrations (including photographs, graphs and diagrams) should be referred to as Figures and their position indicated in the text (e.g., Figure 3). The captions of all figures should be submitted on a separate sheet, should include keys to symbols, and should make interpretation possible without reference to the text. Figures should ideally be professionally drawn and designed with the format of the journal in mind and should be capable of reduction.

Color Figures: Avoid the use of color and tints for purely aesthetic reasons. Any figure submitted as a color original will appear in colour in the journal's online edition free of charge and can be downloaded. Paper copy color reproduction will only be considered on condition that authors contribute to the associated costs. Charges are: US\$500 per article, a 15% service fee will be applied by Rightslink. (Color costs will be waived for invited articles).

Supplemental digital content

The space available in the printed journal is restricted and authors are therefore encouraged to be as concise as possible in their writing and to consider having supplementary material for publication only in the online version of the journal. Supplementary material must be important ancillary information that is relevant to the parent article but is not essential in the printed journal.

Supplemental digital content may enhance an article's text and may include text documents, questionnaires, tables, figures, graphics, illustrations, audio and DVD/video material, research protocols and mathematical calculations. Authors reporting results from a questionnaire survey should include a copy of the questionnaire used (original language and/or English translation) together with the manuscript (see supplementary material above), unless the questionnaire is in common use and published before, when a reference will suffice. Do not include supplementary material within the main manuscript file, but upload as separate file(s).

Cite all supplemental digital content consecutively in the text. Citations should be clearly labelled as "Supplemental Digital Content" on submission. Include a sequential number (S1, S2, S3 etc.) and provide a brief description of the supplemental content. Provide a legend on supplemental digital content at the end of the text. List each legend in the order in which the material is cited in the text.

The legends must be numbered to match the citations from the text. Include a title and a brief summary of the content. For audio and video files, also include the author name, videographer, participants, length (minutes), and size (MB).

To ensure quality for those viewing supplemental digital content, the publisher suggests that authors submit supplemental digital files no larger than 10 MB each. Follow these guidelines for formatting files:

- ☐ Documents, graphs, and tables may be presented in any format.
- ☐ Figures, graphics, and illustrations should be submitted with the following file extensions: .tif, .eps, .ppt, .jpg, .pdf.
- ☐ Audio files should be submitted with the following file extensions: .mp3, .wma.
- ☐ Video files should be submitted with the following file extensions: .wmv, .mov, .qt, .mpg, .mpeg, .mp4. Video files should also be formatted with a 320 _ 240 pixel minimum screen size.

Just Accepted Publication

To provide our readers with the most rapid access to accepted articles, this Journal pre-publishes accepted articles shortly after acceptance as “Just Accepted” articles. Just Accepted articles have undergone full scientific review but none of the editorial preparation, such as copyediting, typesetting, and proofreading, which is mandatory for all articles published in the regular issues of the journal. Publication in the Just Accepted area will send a citation record to PubMed.

Once an article has been copyedited and the final corrected proofs have been typeset, it will move to our Early Online section if it has not yet been assigned to an issue.

Please note that although Just Accepted and Early Online articles do not have all bibliographic details available yet (for example pagination), they can still be cited using the year of online availability and the DOI number as follows: Author(s), Article Title, Journal (Year), DOI.

Authors are requested to make sure that they follow all instructions provided to them at the revision and acceptance stages, to help us to publish their manuscripts as quickly as possible after acceptance. For example authors should remember to delete any redundant manuscript files before completing the submission of your revised manuscript, making sure that all figures/tables and references are correctly supplied and formatted according to the journal style, and most importantly authors should return a signed copy of the Copyright Transfer Agreement to the Publisher as soon as possible to avoid any delays in the online publication of their accepted manuscript as a Just Accepted article*.

*Please note that whilst Informa Healthcare will strive to post all accepted manuscripts within two days of their acceptance, this is contingent on the timely receipt of signed Copyright Transfer Agreements from the author(s) and there may be editorial considerations, such as a planned press release or the need to coordinate posting of related articles, which may preclude or delay Just Accepted posting.

Disclaimer

Informa Healthcare makes every effort to ensure the accuracy of all the information (the “Content”) contained in its publications. However, Informa Healthcare and its agents and licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content and disclaim all such representations and warranties whether express or implied to the maximum extent permitted by law. Any views expressed in this publication are the views of the authors and are not the views of Informa Healthcare.